



**UNIVERSITI PUTRA MALAYSIA**

**EFFECTS OF CHITOSAN IN EXTENDING THE VASE LIFE OF CUT  
CHRYSANTHEMUM (DENDRANTHEMA MORIFOLIUM RAMAT)  
FLOWERS**

**EDDY AZIAN.**

**FS 2006 27**

**EFFECTS OF CHITOSAN IN EXTENDING  
THE VASE LIFE OF CUT CHRYSANTHEMUM  
(*DENDRANTHEMA MORIFOLIUM* RAMAT) FLOWERS**

**By**

**EDDY AZIAN**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfilment of the Requirements for the Degree of Master of Science**

**March 2006**



*..... Ya Allah, ya Tuhan, aku berlindung kepadaMu dari ilmu yang tidak berguna, dari hati yang tidak tunduk, dari nafsu yang tidak puas, dan dari permohonan yang tidak diterima. (H.R. Muslim)*

#### **SPECIAL DEDICATIONS**

This Thesis is dedicated to

my beloved parents, Zahari Nur and Hilalliah

my beloved parents in law, Supriatin and E. Muchtar

my beloved wife, Neni Mulyani, whose patience, supports  
and companionship has facilitated my work,

my daughter, Miza Hilmiya

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in  
fulfilment of the requirement for the degree of Master of Science

**EFFECTS OF CHITOSAN IN EXTENDING THE VASE LIFE OF CUT  
CHRYSANTHEMUM (*DENDRANTHEMA MORIFOLIUM* RAMAT)  
FLOWERS**

By

**EDDY AZIAN**

**March 2006**

**Chairman: Associate Professor Mohamad Zaki Ab. Rahman, PhD**

**Faculty: Science**

The effect of the isolated bacteria strains on the vase life of cut flowers of Chrysanthemum (*Dendranthema morifolium* Ramat) was investigated. The bacteria strains were isolated from the cut stem end and vase water of cut chrysanthemum flowers. The longevity of vase life and total water uptake of cut chrysanthemum was reduced by placing the stems in a suspension of *Pantoea agglomerans* and *Burkholderia multivorans* at  $10^8$  CFU mL<sup>-1</sup>.

The study was also conducted to look at the effects of chitosan in extending the vase life of cut chrysanthemum. Chitosan was applied as pulse treatment. Cut flowers were treated with 1) chitosan at 0, 25, 50, 75 and 100 mg L<sup>-1</sup>, 2) chitosan at concentration 1) supplemented with 2% sucrose, 3) chitosan at concentration 1) supplemented with 1.5 mM Ca(NO<sub>3</sub>)<sub>2</sub> and 4) chitosan at concentration 1) supplemented with 1.5 mM Ca(NO<sub>3</sub>)<sub>2</sub> and 2% sucrose. The experiment was

carried out in the Postharvest Laboratory with the environmental conditions,  $25 \pm 2^{\circ}\text{C}$ , 70% RH and  $15 \mu\text{mol m}^{-2} \text{s}^{-1}$  light intensity from cool-white fluorescent lamps for 12 hours. Compared to those held in control, all treatments extended the vase life. Treatment  $75 \text{ mg L}^{-1}$  chitosan supplemented with  $1.5 \text{ mM Ca}(\text{NO}_3)_2$  was the most effective in extending the vase life.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia  
sebagai memenuhi keperluan untuk ijazah Master Sains

**KAJIAN TERHADAP PENGGUNAAN KITOSAN UNTUK  
MELANJUTKAN HAYAT JAMBANGAN BUNGA KERATAN KEKWA  
(*DENDRANTHEMA MORIFOLIUM* RAMAT)**

Oleh

**EDDY AZIAN**

**Mac 2006**

**Pengerusi: Profesor Madya Mohamad Zaki Ab. Rahman, PhD**

**Fakulti: Sains**

Kesan strain bakteria yang diasingkan terhadap hayat jambangan bunga keratan kekwa (*Dendranthema morifolium* Ramat) telah dikaji. Strain bakteria diasingkan daripada pangkal batang dan cecair pasu bunga keratan kekwa. Tempoh hayat jambangan dan jumlah serapan air bunga keratan kekwa berkurangan apabila batang bunga kekwa diletakkan dalam suspensi *Pantoea agglomerans* dan *Burkholderia multivorans* pada populasi  $10^8$  CFU mL<sup>-1</sup>.

Satu kajian lagi juga dilakukan untuk melihat kesan kitosan dalam memanjangkan hayat jambangan bunga keratan kekwa. Kitosan diperlakukan sebagai rawatan *pulsing*. Bunga keratan kekwa dirawat dengan 1) kitosan pada kepekatan 0, 25, 50, 75 dan 100 mg L<sup>-1</sup>, 2) kitosan pada kepekatan 1) dengan tambahan 2% sukrosa, 3) kitosan pada kepekatan 1) dengan tambahan 1.5 mM Ca(NO<sub>3</sub>)<sub>2</sub> dan 4) kitosan pada kepekatan 1) dengan tambahn 1.5 mM Ca(NO<sub>3</sub>)<sub>2</sub> dan 2% sukrosa. Kajian dilakukan dalam makmal pascatuai dengan suhu persekitaran  $25 \pm 2^\circ\text{C}$ ,

70% RH and  $15 \mu\text{mol m}^{-2} \text{s}^{-1}$  cahaya lampu putih selama 12 jam sehari. Jika dibandingkan dengan kawalan, semua rawatan memanjangkan hayat jambangan. Dengan rawatan  $75 \text{ mg L}^{-1}$  kitosan bersama  $1.5 \text{ mM Ca(NO}_3)_2$  memberikan pemanjangan hayat jambangan yang terbaik.

## ACKNOWLEDGEMENTS

In the name of Allah the Beneficent and the Compassionate. Praise to almighty Allah "S.W.T" for his blessings that enable me to complete this thesis and achieve the academic degree.

I wish to express my deepest gratitude and sincere appreciation to Associate Professor Dr Hj. Mohamad Zaki Ab. Rahman, the chairman of the supervisory committee for his guidance, constant encouragement, invaluable suggestions and generous help throughout the study period and in preparing this manuscript.

My sincere appreciation also goes to the members of my supervisory committee, Assoc. Prof. Dr. Mahmud Tengku Muda Mohamed for giving me the opportunity to work on the project, his interest in progress of the research and constructive discussion for better understanding of the subject. Assoc. Prof. Dr. Kamaruzzaman Sijam for their advice, constructive suggestions and critical reviewing this manuscript.

I am thankful to all of my friends and colleagues, especially for their help and moral encouragement. A very special thank you Mr. Adiwirman, Mr. Bambang Prasetyo, Mr. Anwar and Mr. Teguh Prasetyo. I also acknowledge the help of technical staffs from the Faculty of Agriculture, UPM, especially Mr. Nur, Mr. Azhar, Mr. Mohamad Daud, Mrs. Junainah and Mr. Zawawi.





I am grateful to my family, particularly my mother and father, my parent in laws, sisters and brothers, also to my beloved wife, Neni Mulyani Muchtar, SP and my beloved daughter Miza Hilmiya for their immense help, encouragement, inspiration, patience, unfailing support and sacrifice.

I acknowledge the Universiti Putra Malaysia and the Ministry of Science, Technology and Environment (MOSTE) Malaysia for the financial support during my study, through project IRPA grant No. 01-02-04-0061-EA001, that was awarded to my co-supervisor, Assoc. Prof. Dr. Mahmud Tengku Muda Mohamed, and through special allowance scheme.

I certify that an Examination Committee has met on 1<sup>st</sup> March 2006 to conduct the final examination of Eddy Azian on his Master of Science thesis entitled “Effects of Chitosan in Extending the Vase Life of Cut Chrysanthemum (*Dendranthema morifolium* Ramat) Flowers” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulation 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**Anuar Kassim, PhD**

Professor  
Faculty of Science  
Universiti Putra Malaysia  
(Chairman)

**Phebe Ding, PhD**

Lecturer  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Internal Examiner)

**Mohd. Aspollah Hj. Md. Sukari, PhD**

Associate Professor  
Faculty of Science  
Universiti Putra Malaysia  
(Internal Examiner)

**Md. Pauzi Abdullah, PhD**

Professor  
Faculty of Science and Technology  
Universiti Kebangsaan Malaysia  
(External Examiner)



**HASANAH MOHD. GAZALI, PhD**

Professor/Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 26 APR 2006

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for degree of Master of Science. The members of the Supervisory Committee are as follows:

**Mohamad Zaki Ab. Rahman, PhD**

Associate Professor  
Faculty of Science  
Universiti Putra Malaysia  
(Chairman)

**Mahmud Tengku Muda Mohamed, PhD**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**Kamaruzzaman Sijam, PhD**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)



---


**AINI IDERIS, PhD**

Professor/Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 11 MAY 2006

## DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



**EDDY AZIAN**

Date: *20 April 2006*

## TABLE OF CONTENTS

		<b>Page</b>
	<b>DEDICATION</b>	ii
	<b>ABSTRACT</b>	iii
	<b>ABSTRAK</b>	v
	<b>ACKNOWLEDGEMENTS</b>	vii
	<b>APPROVAL</b>	ix
	<b>DECLARATION</b>	xi
	<b>LIST OF TABLES</b>	xvii
	<b>LIST OF FIGURES</b>	xviii
	<b>LIST OF ABBREVIATION</b>	xxi
	<b>CHAPTER</b>	
<b>I</b>	<b>INTRODUCTION</b>	1
<b>II</b>	<b>LITERATURE REVIEW</b>	5
	2.1 Chitosan	5
	2.1.1 Introduction	5
	2.1.2 Preparation of chitin and chitosan	7
	2.1.3 Present chitin and chitosan research in Asia	9
	2.1.4 Antimicrobial activities of chitosan	9
	2.1.5 Chitosan in the edible film industry	14
	2.2 Chrysanthemum	16
	2.2.1 Botanical description	16
	2.3 Factors affecting vase life	17
	2.3.1 Vascular blockage	17
	2.3.1.1 Physiological factors	18
	2.3.1.2 Microbiological factors	19
	2.3.2 Cavitation and embolisation	20
	2.3.3 Ethylene and cut flower	21
	2.4 Preservative solution	22
	2.4.1 Water	23
	2.4.2 Organic acids and salts; and antioxidants	24
	2.4.3 Surfactants and wetting agents	25
	2.4.4 Germicides	25
	2.4.5 Sugars	26
<b>III</b>	<b>ISOLATION OF MICROORGANISMS FROM VASE SOLUTION OF CHRYSANTHEMUMS AND EFFECT OF DIFFERENT ISOLATED BACTERIAL STRAINS ON THE VASE LIFE</b>	27
	3.1 Introduction	27
	3.2 Material and methods	29



3.2.1	Isolation and purification of microorganisms in stem end and vase water	30
3.2.1.1	Preparation of cut flowers	30
3.2.1.2	Identification of microorganisms	31
3.2.2	Effect of isolated bacteria strains on the vase life of cut chrysanthemum flowers	33
3.2.2.1	Preparation of vase solution inoculation	33
3.2.2.2	Preparation of cut flowers	33
3.2.2.3	Determination of water uptake	34
3.2.2.4	Determination of fresh weight	34
3.2.2.5	Measurement of flower diameter	35
3.2.2.6	Vase life observations	35
3.2.3	Scanning electron microscopy (SEM) specimen preparation	35
3.2.4	Experimental design and statistical analysis	36
3.3	Results	37
3.3.1	The fungus in stems and vase solution of cut chrysanthemum	37
3.3.2	The bacterial in stems and vase solution of cut chrysanthemum	37
3.3.3	Effect of bacterial strains in the vase water on vase life of cut chrysanthemums	39
3.3.4	Scanning electron microscopy (SEM)	43
3.4	Discussion	44
3.5	Conclusion	46
IV	<b>VASE LIFE OF CUT CHRYSANTHEMUM (<i>DENDRANTHEMA MORIFOLIUM</i> RAMAT) AS AFFECTED BY CHITOSAN SUPPLEMENTS</b>	47
4.1	Introduction	47
4.2	Material and methods	49
4.2.1	Preparation of chitosan from shell of shrimp	49
4.2.2.1	Deproteinization	49
4.2.2.2	Demineralization	49
4.2.2.3	Deacetylation	50
4.2.2	Preparation of chitosan solution	50
4.2.3	Preparation of cut flowers	51
4.2.4	Pulse treatments	51
4.2.5	Treatment	52
4.2.5.1	Determination of water uptake	53
4.2.5.2	Determination of fresh weight	54
4.2.5.3	Measurement of flower diameter	54
4.2.5.4	Vase life observations	55

4.3	Experimental design and statistical analysis	55
4.4	Results	56
4.4.1	The effect of pulse treatment with chitosan on quality of cut chrysanthemum flowers	56
4.4.2	The effect of pulsing in different concentrations of chitosan supplemented with sucrose	61
4.4.3	The effect of pulsing in different concentrations chitosan supplemented with $\text{Ca}(\text{NO}_3)_2$	66
5.4.4	The effect of pulsing in different concentration of chitosan supplemented with sucrose and $\text{Ca}(\text{NO}_3)_2$	71
4.5	Discussion	76
4.6	Conclusion	76
<b>V</b>	<b>GENERAL DISCUSSION AND CONCLUSIONS</b>	<b>77</b>
	<b>BIBLIOGRAPHY</b>	<b>81</b>
	<b>APPENDICES</b>	<b>90</b>
	<b>BIODATA OF THE AUTHOR</b>	<b>98</b>

## LIST OF TABLES

Table		Page
1	Identification of bacteria strains isolated from the stem end and vase solution after 10 days vase life observation of cut chrysanthemum flowers.	37
2	The effects of chitosan at various concentrations on total water uptake (g) during seven days of observation. * Note: Viscosity measured at 0 day.	57
3	The effects of chitosan addition in 2% sucrose pulse treatment on the total water uptake (g) during seven days of observation. *Note: Viscosity measured at 0 day.	61
4	Effect of pulse treatment with various concentrations of chitosan supplemented with 1.5 mM $\text{Ca}(\text{NO}_3)_2$ on the total water uptake (g) during seven days of observation. * Note: Viscosity measured at 0 day.	66
5	Effect of pulse treatment with various concentrations of chitosan supplemented with 2% sucrose and 1.5 mM $\text{Ca}(\text{NO}_3)_2$ on the total water uptake (g) during seven days of observation. *Note: Viscosity measured at 0 day.	71



## LIST OF FIGURES

Figure	Page
1	7
2	8
3	38
4	39
5	40
6	41
7	42
8	43
9	52
10	56
11	58

Note: Means followed by the same letter among the treatments are not significantly different at ( $p<0.05$ ) as determined by DMRT.

12	The effect of pulse treatment with various concentrations of chitosan on percentage relative fresh weight of cut chrysanthemum during 14 days of observation.	58
13	The effect of pulse treatment with various concentrations of chitosan on diameter of flower. Note: Means followed by the same letter among the treatments are not significantly different at ( $p<0.05$ ) as determined by DMRT.	59
14	The effects of pulse treatment with various concentrations of chitosan supplemented with 2% sucrose on the vase life of cut chrysanthemum flowers. Note: Means followed by the same letter among the treatments are not significantly different at ( $p<0.05$ ) as determined by DMRT.	60
15	The effect of pulse treatment with various concentrations of chitosan supplemented with 2% sucrose on fresh weight of cut chrysanthemum flowers. Note: Means followed by the same letter among the treatments are not significantly different at ( $p<0.05$ ) as determined by DMRT.	62
16	The effect of pulse treatment with various concentrations of chitosan supplemented with 2% sucrose on relative fresh weight of cut chrysanthemum flowers during 14 days of observation.	63
17	The effect of pulse treatment with various concentrations of chitosan supplemented with 2% sucrose on diameter of flower. Note: Means followed by the same letter among the treatments are not significantly different at ( $p<0.05$ ) as determined by DMRT.	64
18	The effect of pulse treatment with various concentrations of chitosan supplemented with 1.5 mM $\text{Ca}(\text{NO}_3)_2$ on the vase life of cut chrysanthemum flowers. Note: Means followed by the same letter among the treatments are not significantly different at ( $p<0.05$ ) as determined by DMRT.	65
19	The effect of pulse treatment with various concentrations of chitosan supplemented with 1.5 mM $\text{Ca}(\text{NO}_3)_2$ on fresh weight of cut chrysanthemum flowers. Note: Means followed by the same letter among the treatments are not significantly different at ( $p<0.05$ ) as determined by DMRT.	67

20	The effect of pulse treatment with various concentrations of chitosan supplemented with 1.5 mM $\text{Ca}(\text{NO}_3)_2$ on percentage relative fresh weight of cut chrysanthemum during 14 days of observation.	68
21	The effect of pulse treatment with various concentrations of chitosan supplemented with 1.5 mM $\text{Ca}(\text{NO}_3)_2$ on diameter of flower. Note: Means followed by the same letter among the treatments are not significantly different at ( $p < 0.05$ ) as determined by DMRT.	69
22	The effect of chitosan at different concentrations supplemented with 2% sucrose + 1.5 mM $\text{Ca}(\text{NO}_3)_2$ in pulse treatment on the vase life of cut chrysanthemum flowers. Note: Means followed by the same letter among the treatments are not significantly different at ( $p < 0.05$ ) as determined by DMRT.	70
23	The effect of pulse treatment with various concentrations of chitosan supplemented with 2% sucrose and 1.5 mM $\text{Ca}(\text{NO}_3)_2$ on fresh weight of cut chrysanthemum flowers. Note: Means followed by the same letter among the treatments are not significantly different at ( $p < 0.05$ ) as determined by DMRT.	72
24	The effect of pulse treatment with various concentrations of chitosan supplemented with 2% sucrose and 1.5 mM $\text{Ca}(\text{NO}_3)_2$ on percentage relative fresh weight of cut chrysanthemum during 14 days of observation.	73
25	The effect of pulse treatment with various concentrations of chitosan supplemented with 2% sucrose and 1.5 mM $\text{Ca}(\text{NO}_3)_2$ on diameter of flower. Note: Means followed by the same letter among the treatments are not significantly different at ( $p < 0.05$ ) as determined by DMRT.	74

## LIST OF ABBREVIATIONS

AIB	aminoisobutyric acid
ANOVA	analysis of variance
atm	atmosphere
BCDMH	1-bromo-3-chloro-5,5-dimethylhydantoin
°C	degree centigrade (celcius)
CFU	colony forming unit
cm	centimetre
CRD	completely randomized design
DICA	dichloroisocyanuric acid
DMRT	Duncan's multiple range test
e.g.	for example
<i>et al.</i>	et alteri
g	gram
GlcN	<i>N</i> -acetyl- $\beta$ -D-glucosamine
GN	gram negative
ha	hectare
HQC	hydroxyquinoline citrate
klx	kilolux
KOH	potassium hydroxide
L	litre
$\mu$	micro ( $10^{-6}$ )
M	molar

m	milli ( $10^{-3}$ )
MIC	minimum inhibitory concentration
MT	metric ton
MW	molecular weights
ppm	part per million
RH	relative humidity
s	second
SEM	scanning electron microscopy
ST	simulated transport
STS	silver thiosulphate
v/v	volume / volume
w/v	weight / volume
w/w	weight / weight

## CHAPTER I

### INTRODUCTION

The vase life longevity is one of the most important quality factors for cut flowers (Pun *et al.*, 1999; Ichimura *et al.*, 2002). Most of cut flowers show leaf wilting shortly after being placed in the vase solution. Cut flowers are kept at room temperature in houses for decoration and end up dry mostly through water loss (Anjum *et al.*, 2001). This water stress develops soon after placing the flowers in water and this often cause shortening of vase life. Unrestricted water uptake through the xylem conduits is of paramount importance in providing an adequate supply of water to transpire cut stems. According to Havelly and Mayak (1981) and van Doorn (1997), the main reason for water stress during the vase period is resistance to water flow which develops in the stems.

According to van Doorn *et al.* (2002) the water uptake during vase life of flowers can be shortened, due to development of an occlusion in the xylem conduits, leading to premature wilting of the leaves and flowers. There are several causes of occlusion, such as microbial contamination in the water and microbial growth in the stems, and a physiological response for cutting.

The presence of bacteria in vase water greatly influences the longevity of cut flowers. This was observed in cut carnation (van Doorn *et al.*, 1995), roses (Lineberger and Steponkus, 1976; de Witte and van Doorn, 1988; van Doorn and

Perik, 1990; Florack *et al.*, 1996), and *Adiantum raddianum* frond (van Doorn *et al.*, 1991). Besides that, the presence of microbial in vase water can also cause release of toxic metabolites and/or enzymes, and evolve damaging levels of ethylene (Gilman and Steponkus, 1972; van Doorn and Perik, 1990; Jones and Hill, 1993).

The use of floral preservative is the most economical and practicle method of extending the postharvest life of cut flowers. Several preservative such as silver thiosulphate (STS), silver nitrate ( $\text{AgNO}_3$ ), and aluminum sulphate have physiological effects on vase life and have been reported to promote flowers fresh weight increase, reduced microbial numbers, and inhibited the effects of ethylene. However, chemical control program causes environmental problems, such as, an increasing number of fungicides-resistant strains of postharvest pathogens, and a number of commonly used fungicides are under review due to health risk concerns (El Ghaouth *et al.*, 1991; Florack *et al.*, 1996). Thus, exploring ways to replace these chemical compounds devoid of any heavy metal (Onozaki *et al.*, 1998) by natural products with antibacterial properties or to intensify the natural defences of the tissue to control decay and prolong storage life is receiving more and more attention (Florack *et al.*, 1996; Zhang and Quantick, 1998).

Chitosan is a linear polymer composed of  $\beta$ -1,4-linked glucosamine (GlcN) with various degrees of N-acetylated GlcN residues, is a deacetylated derivative of chitin extracted from an abundant source of shellfish exoskeletons. The amino

polysaccharide and its derivatives have many diverse applications in agriculture, medicine and cosmetics (Kittur *et al.*, 2003).

The use of a natural substance such as chitosan has been considered as a valid alternative to synthetic fungicides (Romanazzi *et al.*, 2003) because the polymer which is naturally occur should not cause health and pollution problems. The unique positively charged chitosan in acidic solution has potential uses in agriculture such as a carrier for the controlled release of fungicides and fertilizers. According to Jung *et al.* (1999), antifungal or antimicrobial activities of chitosan are believed to originate from the polycationic nature of chitosan that can be bind with anionic sites in proteins, thus resulting in selective antimicrobial activities towards fungi or bacteria.

Several recent studies have shown that chitosan has antibacterial and antifungal characteristics. Plant seeds soaked in aqueous solution of chitosan can prevent microbial infections and increased plant production. Ohta *et al.* (1999) has reported that chitosan promote growth of *Eustoma grandiflorum* seedlings and flowering.

Chitosan is used as spray or coating of fresh fruits and vegetables. Coating fruit with chitosan prolongs the storage life and better control of decay in tomatoes, and strawberry fruits (El Ghaouth *et al.*, 1992a and b), litchi, peach, Japanese pear and kiwifruit (Zhang and Quantick, 1997; Du *et al.*, 1997). Makino and Hirata (1997) reported that the biodegradable laminate chitosan-cellulose was able to use as



packaging materials for storage of fresh vegetables (head lettuce, cut broccoli, whole broccoli, tomatoes, sweet corn and blueberries). Recently, Reddy *et al.* (2000) found that spray application of chitosan affected quality of strawberry fruits and having potential as an anti-transpirant of pepper (Bittelli *et al.*, 2001).

The use of edible films and coating to extend shelf life and improve the quality of fresh, frozen and food industry has been examined during the past few years (Kester and Fennema, 1986). The film-forming qualities of chitosan have been found to be dependent on the homogeneity of the bulk material, the degree of acetylation and acid degradation during dissolution. A composite film derived from chitosan have been developed with qualities such as good wet strength, excellent oxygen barrier properties and biodegradability (Hosokawa *et al.*, 1990; Conca and Yang, 1993; Caner *et al.*, 1998).

Most of the studies previously carried out only indicated that chitosan have been applied in postharvest treatments and preharvest application to extend storage life of fruits and vegetables. However, information of the use as cut flower preservation are still lacking. Therefore, the objectives of this study were:

1. To evaluate the role of bacterial strains in the vase life of cut chrysanthemum.
2. To evaluate the cut chrysanthemum flower quality by using chitosan supplements.